

Late Safety, Efficacy, and Cost-Effectiveness of a Zotarolimus-Eluting Stent Compared With a Paclitaxel-Eluting Stent in Patients With De Novo Coronary Lesions

2-Year Follow-Up From the ENDEAVOR IV Trial (Randomized, Controlled Trial of the Medtronic Endeavor Drug [ABT-578] Eluting Coronary Stent System Versus the Taxus Paclitaxel-Eluting Coronary Stent System in De Novo Native Coronary Artery Lesions)

Martin B. Leon, MD,* David E. Kandzari, MD,† Eric L. Eisenstein, DBA,‡ Kevin J. Anstrom, PhD,‡ Laura Mauri, MD, MSc,§ Donald E. Cutlip, MD,|| Eugenia Nikolsky, MD,* Charles O'Shaughnessy, MD,¶ Paul A. Overlie, MD,# Ajay J. Kirtane, MD,* Brent T. McLaurin, MD,** Stuart L. Solomon, MD,†† John S. Douglas Jr, MD,‡‡ Jeffrey J. Popma, MD,|| for the ENDEAVOR IV Investigators

New York, New York; La Jolla, California; Durham, North Carolina; Boston, Massachusetts; Elyria, Ohio; Lubbock, and Houston, Texas; Anderson, South Carolina; and Atlanta, Georgia

Objectives The aim of this study was to assess, after 2 years of follow-up, the safety, efficacy, and cost-effectiveness of a zotarolimus-eluting stent (ZES) compared with a paclitaxel-eluting stent (PES) in patients with native coronary lesions.

Background Early drug-eluting stents were associated with a small but significant incidence of very late stent thrombosis (VLST), occurring >1 year after the index procedure. The ZES has shown encouraging results in clinical trials.

Methods The ENDEAVOR IV trial (Randomized, Controlled Trial of the Medtronic Endeavor Drug [ABT-578] Eluting Coronary Stent System Versus the Taxus Paclitaxel-Eluting Coronary Stent System in De Novo Native Coronary Artery Lesions), a randomized (1:1), single-blind, controlled trial (n = 1,548) compared ZES versus PES in patients with single de novo coronary lesions. Two-year follow-up was obtained in 96.0% of ZES and 95.4% of PES patients. The primary end point was target vessel failure (TVF), and safety end points included Academic Research Consortium-defined stent thrombosis. Economic end points analyzed included quality-adjusted survival, medical costs, and relative cost-effectiveness of ZES and PES.

Results The TVF at 2 years was similar in ZES and PES patients (11.1% vs. 13.1%, p = 0.232). There were fewer myocardial infarctions (MIs) in ZES patients (p = 0.022), due to fewer periprocedural non-Q-wave MIs and fewer late MIs between 1 and 2 years. Late MIs were associated with increased VLST (PES: 6 vs. ZES: 1; p = 0.069). Target lesion revascularization was similar comparing ZES with PES (5.9% vs. 4.6%; p = 0.295), especially in patients without planned angiographic follow-up (5.2% vs. 4.9%; p = 0.896). The cost-effectiveness of ZES and PES was similar.

Conclusions After 2 years of follow-up, ZES demonstrated efficacy and cost-effectiveness comparable to PES, with fewer MIs and a trend toward less VLST. (The ENDEAVOR IV Clinical Trial: A Trial of a Coronary Stent System in Coronary Artery Lesions; [NCT00217269](#)) (J Am Coll Cardiol Intv 2009;2:1208–18) © 2009 by the American College of Cardiology Foundation

From *Columbia University Medical Center and the Cardiovascular Research Foundation, New York, New York; †Scripps Clinic, La Jolla, California; ‡Duke Medical Center, Durham, North Carolina; §Harvard Clinical Research Institute, Brigham and Women's Hospital, and ||Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts; ¶Elyria

Studies with Food and Drug Administration-approved drug-eluting stents (DES) with bio-stable polymers to deliver potent antiproliferative agents (sirolimus, paclitaxel, zotarolimus, and everolimus) have demonstrated an improvement in angiographic and clinical efficacy and comparable safety when compared with bare-metal stents (BMS) in randomized clinical trials after 1-year follow-up (1–6).

See page 1236

However, unlike BMS, early versions of DES (sirolimus-eluting stent [SES] and paclitaxel-eluting stent [PES]) were associated with a disturbing complication, very late stent thrombosis (VLST), which occurred after the first year of DES implantation (7–9) and had previously only been observed after coronary vascular brachytherapy procedures (10,11). Although VLST after DES is a rare event, occurring at a frequency varying from 0.2% to 0.6%/year (8), the clinical significance cannot be discounted, because almost all patients suffer myocardial infarctions (MIs) or death. The VLST hazard might be constant for several years (12), and there is an increased sensitivity to antiplatelet therapy withdrawal (13–15). Thus, clinical practice must be modulated to account for possible VLST after DES, which has resulted in more selective DES use and prolonged dual antiplatelet therapy considerations (16,17).

The design characteristics of the zotarolimus-eluting stent (ZES) were intended to optimize patient safety. Experimental results in animal models have indicated that ZES has improved healing responses and restored endothelial function compared with SES and PES, more closely resembling a BMS (18–20). Surrogate safety end points in patients, with intravascular ultrasound, angiography, and optical coherence tomography, have also shown beneficial ZES responses compared with earlier SES and PES (21–24), and a pooled analysis with long-term follow-up of ZES-treated patients has indicated a frequency of VLST <1%, which is similar to results after BMS treatment (25,26). Initial findings from the ENDEAVOR IV trial (Randomized, Controlled Trial of the Medtronic Endeavor Drug [ABT-578] Eluting Coronary Stent System Versus the Taxus Paclitaxel-Eluting Cor-

onary Stent System in De Novo Native Coronary Artery Lesions) involving 1,548 patients with de novo native coronary lesions have demonstrated similar clinical safety and efficacy of ZES compared with PES during the first year of follow-up (5). The purpose of this report is to analyze clinical safety, efficacy, and cost effectiveness with extended 2-year follow-up in the ENDEAVOR IV trial.

Methods

Study design and patient population. The ENDEAVOR IV trial was a prospective, multicenter, single-blinded, randomized, controlled clinical trial that compared clinical and angiographic outcomes between patients treated with ZES and patients treated with PES. Consecutive adult patients with clinical evidence of ischemic coronary disease or a positive functional study were enrolled at 80 centers in the U.S. The institutional review board at each site approved the protocol, and each eligible patient provided written, informed consent before the index procedure.

Description of the key inclusion and exclusion criteria, study devices, and procedural-related details has been previously reported (5). After the interventional procedure, patients were evaluated at 30 days, 6, 9, and 12 months and yearly thereafter up to 5 years after the procedure. Patients in the angiographic and intravascular ultrasound subgroups were assessed at 8 months after the procedure. Dual antiplatelet therapy (aspirin and clopidogrel) was given for 6 months to all patients per protocol and continued thereafter, at the discretion of the managing physician.

Abbreviations and Acronyms

ARC = Academic Research Consortium

BMS = bare-metal stent(s)

DES = drug-eluting stent(s)

DRG = diagnosis-related group

MACE = major adverse cardiac events

MI = myocardial infarction

PES = paclitaxel-eluting stent(s)

QALY = quality-adjusted life year

ST = stent thrombosis

TLR = target lesion revascularization

TVF = target vessel failure

VLST = very late stent thrombosis

ZES = zotarolimus-eluting stent(s)

Memorial Hospital, Elyria, Ohio; #Lubbock Heart Hospital, Lubbock, Texas; **Anderson Medical Center, Anderson, South Carolina; ††Methodist Hospital, Houston, Texas; and the ‡Emory University, Atlanta, Georgia. The ENDEAVOR IV clinical trial was funded by Medtronic CardioVascular, Santa Rosa, California. This work was supported by Medtronic CardioVascular, Santa Rosa, California. Dr. Leon has served as a consultant to Abbott Vascular, Boston Scientific, Cordis, Medtronic, Inc., and Volcano Corporation. Dr. Kandzari has received research/grant support and consulting honoraria from Medtronic, Inc. and Cordis Corporation. Eric L. Eisenstein has received research or grant support from Medtronic, Inc. and Eli Lilly and Company. Dr. Anstrom has received research support from AstraZeneca, Bristol-Myers Squibb, Eli Lilly and Co., and Medtronic, Inc., and has served as a consultant for Johnson & Johnson and Pfizer. Dr. Mauri has served as a consultant to Abbott Vascular, Boston Scientific, Cordis, and Medtronic, Inc. Dr. O'Shaughnessy has received research grants from Medtronic, Inc. Dr. Kirtane has served as a consultant and speaker for Medtronic CardioVascular and Abbott Vascular. Dr. Douglas has served as a clinical investigator for Abbott Vascular, Boston Scientific, Cordis Corporation, Medtronic, and St. Jude Medical. Dr. Popma has received research grants from Abbott Vascular, Biosensors, Boston Scientific, Cordis, ev3, and Medtronic; he has served as a consultant for Abbott Vascular, Boston Scientific, Cordis, Lilly, and Medtronic, and a speaker for Boston Scientific, Bristol-Myers Squibb, Cordis, Lilly, Medicines Co, Medtronic, Pfizer and Sanofi. David O. Williams, MD, served as Guest Editor for this article.

Manuscript received July 31, 2009; revised manuscript received October 13, 2009, accepted October 16, 2009.

Table 1. Baseline Clinical and Angiographic Characteristics of the Study Population

| | ZES | PES | p Value |
|--|---------------------|---------------------|---------|
| Patient demographics | | | |
| Age (yrs) | 63.5 ± 11.1 (773) | 63.6 ± 11.0 (775) | 0.930 |
| Male | 66.9% (517/773) | 68.5% (531/775) | 0.514 |
| Diabetes | 31.2% (241/773) | 30.5% (236/775) | 0.783 |
| Hypertension | 79.4% (614/773) | 82.6% (640/775) | 0.120 |
| Hyperlipidemia | 81.4% (629/773) | 84.8% (657/775) | 0.078 |
| History of smoking | 62.6% (479/765) | 60.4% (462/765) | 0.401 |
| Prior MI | 21.1% (161/764) | 23.2% (176/759) | 0.324 |
| Prior percutaneous coronary intervention | 28.2% (218/773) | 29.5% (229/775) | 0.575 |
| Prior coronary bypass surgery | 9.8% (76/773) | 8.4% (65/775) | 0.332 |
| Angina | | | 0.367 |
| Stable | 45.6% (281/616) | 47.9% (292/609) | |
| Unstable | 51.6% (318/616) | 49.9% (304/609) | |
| MI | 2.8% (17/616) | 2.1% (13/609) | |
| CCS class III or IV | 50.3% (309/614) | 47.9% (292/610) | 0.392 |
| Angiographic characteristics | | | |
| Target vessel | | | 0.791 |
| Left anterior descending | 42.2% (326/772) | 41.5% (321/774) | |
| Left circumflex | 26.9% (208/772) | 26.1% (202/774) | |
| Right coronary | 30.8% (238/772) | 32.4% (251/774) | |
| Type B2/C lesion | 69.6% (537/772) | 70.9% (549/774) | 0.358 |
| Number of diseased, native, major epicardial coronary vessels (>50% stenosed) | | | 0.485 |
| Single | 54.9% (424/772) | 57.2% (443/774) | |
| Double | 28.6% (221/772) | 26.1% (202/774) | |
| Triple | 16.5% (127/772) | 16.7% (129/774) | |
| Left ventricular ejection fraction (%) | 57.3 ± 9.9 (760) | 57.5 ± 10.3 (753) | 0.745 |
| Reference vessel diameter (mm) | 2.73 ± 0.47 (772) | 2.70 ± 0.46 (774) | 0.197 |
| Lesion length (mm) | 13.41 ± 5.67 (771) | 13.80 ± 6.09 (773) | 0.199 |
| Minimal lumen diameter (mm) | 0.96 ± 0.40 (772) | 0.93 ± 0.40 (774) | 0.149 |
| Diameter stenosis (%) | 64.83 ± 13.29 (772) | 65.68 ± 13.10 (774) | 0.204 |
| Values are % (n/total) or mean ± SD (n). | | | |
| CCS = Canadian Cardiovascular Society angina class; MI = myocardial infarction; PES = paclitaxel-eluting stent(s); ZES = zotarolimus-eluting stent(s). | | | |


Clinical end points and definitions. The primary clinical end point in this clinical trial was target vessel failure (TVF), defined as the composite of cardiac death, MI, or clinically driven target vessel revascularization of the treated vessel at 9 months after the procedure. Pre-specified secondary clinical safety and efficacy end points included major adverse cardiac events (MACE), defined as death, MI, or clinically driven target lesion revascularization (TLR) (and individual component end points) and stent thrombosis (ST) at annual follow-up. Myocardial infarction was defined as either: 1) Q-wave MI requiring new pathologic Q waves in 2 or more contiguous electrocardiographic leads and either symptoms consistent with acute myocardial ischemia or elevation of cardiac biomarkers; or 2) non-Q-wave MI requiring elevated creatine kinase >2× the upper laboratory limit with elevated creatine kinase-myocardial band in the absence of pathological Q waves.

Stent thrombosis (both early and late) was adjudicated by a subcommittee of the clinical event committee, blinded to

study stent identity and according to the definitions proposed by the Academic Research Consortium (ARC) (27).

Economic end points and analyses. In addition to clinical outcomes, an economic analysis was performed to determine the relative cost-effectiveness after treatment with ZES or PES. After identification of economic events with MACE- and adverse event-related hospital stays, adverse event system organ class designations were used to distinguish cardiac from noncardiac hospital stays, and Medical Dictionary for Regulatory Activities preferred terms were applied to differentiate types of cardiac hospital stays. Each study subject was assigned an index episode of care and a variable number of follow-up period episodes of care with a method that combined economic event data. Outpatient deaths were the only outpatient events.

Medical costs, length of stay, and quality of life estimates were assigned to episodes of care. The Centers for Medicare and Medicaid Services Medicare Severity Diagnosis Related Grouper was used to associate diagnosis-related groups



(DRGs) to episodes of care (28), and these assignments were then audited by a trained medical record professional. For cardiovascular episodes of care, medical costs and lengths of stay were estimated with Medicare national average payment amounts for 2008 (calculated with an average hospital Medicare base rate of \$4,893) and arithmetic mean lengths of stay by DRG (29,30). For noncardiovascular episodes of care, medical cost and length of stay were assigned with Medicare national average relative weight and arithmetic mean length of stay. The DRG-specific costs for physician services were estimated with published sources and adjusted to 2008 values with the medical care component of the consumer price index (31). A \$2,100 unit cost was assigned to both stent types (data on

file, Medtronic CardioVascular, Santa Rosa, California). Because the type of repeat percutaneous coronary intervention procedure was not identified in the follow-up case report forms for this trial, results were extrapolated from the repeat percutaneous coronary intervention procedures performed at a single participating center (Duke University Medical Center, Durham, North Carolina) during the ENDEAVOR IV study period. These data were used to define the distribution of different types of percutaneous revascularization (32). Previously defined methods were used to assign quality of life weights (33), including a 0.79 quality-adjusted life year (QALY) adjustment for all revascularization procedures, a 0.85 QALY adjustment for coronary artery disease without revascularization, a 0.88 permanent QALY adjustment when subjects experienced a nonfatal MI, and a quality adjusted life day decrement for each day of estimated length of stay during a hospital stay (34).

Number of hospital stays (revascularization, other cardiac, noncardiac, and total), quality-adjusted survival (both undiscounted and discounted at 3%), and medical costs (both undiscounted and discounted at 3%) were computed as 2-year cumulative values by treatment (ZES or PES) with differences, 95% confidence intervals, and p values. Medical costs also were presented by annual time periods. Analyses were performed with generalized linear models on partitioned data with empirical standard errors and an adjustment for censoring. We then performed a comprehensive cost-effectiveness analysis to estimate the incremental medical costs/QALY saved with the use of ZES versus PES during the 2-year follow-up period. This cost-effectiveness ratio was calculated with the nonparametric bootstrap procedure, and results are shown as discounted mean values for the ratio.

Table 2. Clinical Outcomes at 2 Years

| | ZES (n = 742) | PES (n = 739) | p Value |
|---|------------------|------------------|---------|
| Death (all) | 3.1 (23) | 2.6 (19) | 0.639 |
| Cardiac | 1.5 (11) | 1.2 (9) | 0.823 |
| Noncardiac death | 1.6 (12) | 1.4 (10) | 0.831 |
| MI (all) | 2.0 (15) | 4.1 (30) | 0.023 |
| Q-wave | 0.4 (3) | 0.5 (4) | 0.725 |
| Non-Q-wave | 1.6 (12) | 3.5 (26) | 0.022 |
| Death or MI | 5.0 (37) | 6.5 (48) | 0.221 |
| Cardiac death or MI | 3.4 (25) | 5.1 (38) | 0.096 |
| Target lesion revascularization | 5.9 (44) | 4.6 (34) | 0.295 |
| Target vessel revascularization | 8.9 (66) | 9.2 (68) | 0.857 |
| Nontarget lesion, target vessel revascularization | 4.2 (31) | 5.8 (43) | 0.154 |
| Target vessel failure | 11.1 (82) | 13.1 (97) | 0.232 |
| Major adverse cardiac events | 9.8 (73) | 10.0 (74) | 0.931 |

Values are % (n).
Abbreviations as in Table 1.




Figure 2. Cumulative Incidence of TVF and TLR at 2 Years

Hazard ratios (HR) with 95% confidence intervals and cumulative incidence of target vessel failure (TVF) (A) and target lesion revascularization (TLR) (B) at 1 and 2 years are displayed. Abbreviations as in Figure 1.

Statistical methods. The randomized ENDEAVOR IV trial was designed to compare the noninferiority primary end point (TVF at 9 months) between patients treated with ZES and those treated with PES (5). The primary objective of this analysis was to compare clinical safety and effectiveness outcomes between the 2 DES cohorts 2 years after the index revascularization.

Categorical variables were compared by the chi-square or Fisher exact test, as appropriate. Continuous variables are described as means with SD and were compared by unpaired *t* tests. Time-to-event data are reported and displayed as Kaplan-Meier estimates, with comparisons between groups by the log-rank test. All analyses are by

intention-to-treat, with all patients randomized to each study stent included.

All analyses were performed with SAS software (version 8.2 or higher, SAS Institute, Cary, North Carolina). Data collection, clinical event adjudication, and analysis were performed at the Harvard Clinical Research Institute (Boston, Massachusetts). All authors had full access to the database and analysis upon which this manuscript is based.

Results

Clinical outcomes at 2 years. Baseline clinical and angiographic characteristics of the study population were similar




Figure 3. TLR by Angiographic and Clinical Follow-Up to 2 Years

Number and percent TLR for patients with angiographic follow-up and clinical follow-up at 2 years. Abbreviations as in Figure 1.

among ZES and PES patients (Table 1). Of the 1,548 patients enrolled in the ENDEAVOR IV trial, clinical follow-up was completed at 1 and 2 years in 97.5% and 96.0% of ZES patients and in 96.9% and 95.4% of PES patients, respectively, (Fig. 1). The primary study end point, TVF, was 11.1% for ZES compared with 13.1% for PES ($p = 0.232$) (Table 2, Fig. 2A). There were no differences in death, but there were significantly fewer non-Q-wave MIs in ZES versus PES patients (1.6% vs. 3.5%; $p = 0.022$, Table 2). There were no differences in overall clinically driven TLR or target vessel revascularization events comparing ZES with PES at 2 years (Table 2, Fig. 2B). In those patients with angiographic follow-up (18% of the total study population), TLR at 2 years was 9.2% for ZES and 3.1% for PES ($p = 0.045$) (Fig. 3). In those patients without angiographic follow-up, TLR was 5.2% for ZES and 4.9% for PES ($p = 0.896$) (Fig. 3).

ST WITHIN THE FIRST YEAR. Although there were no significant differences in early or late ARC ST during the first year of follow-up among the treatment cohorts, there was a trend toward more frequent ARC-definition (definite or probable) ST with ZES compared with PES (0.9% vs. 0.1%, $p = 0.07$). Of the 7 ZES patients with ARC ST (definite or probable), 3 occurred before 30 days (2 definite and 1 probable) and 4 occurred between 30 days and 6 months (3 definite and 1 probable). In the 3 ZES patients with ARC definite or probable ST before 30 days, 2 were associated with edge dissections and incomplete stent expansion, and 1 had an unplanned surgical procedure. In the 4 ZES patients with ARC definite or probable ST between 30 days and 6 months, 3 patients were no longer taking dual antiplatelet therapy (time of dual antiplatelet therapy cessation to ST was 2 days, 20 days, and 2 months); 1 of which also had an unplanned surgical procedure.

BETWEEN 1 AND 2 YEARS. Academic Research Consortium VLST (definite or probable) occurred in 1 ZES patient and in 6 PES patients ($p = 0.069$) (Tables 3 and 4, Fig. 4). The 6 PES ST events were associated with no deaths and 3

Table 3. Stent Thrombosis Through 2 Years

| Outcome | ZES (n = 742) | PES (n = 739) | p Value |
|--------------------------|------------------|------------------|---------|
| Early (0–30 days) | | | |
| Definite | 0.3 (2) | 0.1 (1) | 1.000 |
| Probable | 0.1 (1) | 0.0 (0) | 1.000 |
| Possible | 0.0 (0) | 0.0 (0) | — |
| Definite/probable | 0.4 (3) | 0.1 (1) | 0.624 |
| Any | 0.4 (3) | 0.1 (1) | 0.624 |
| Late (31–360 days) | | | |
| Definite | 0.4 (3) | 0.0 (0) | 0.249 |
| Probable | 0.1 (1) | 0.0 (0) | 1.000 |
| Possible | 0.4 (3) | 0.4 (3) | 1.000 |
| Definite/probable | 0.5 (4) | 0.0 (0) | 0.124 |
| Any | 0.9 (7) | 0.4 (3) | 0.342 |
| Very late (361–720 days) | | | |
| Definite | 0.1 (1) | 0.7 (5) | 0.124 |
| Probable | 0.0 (0) | 0.1 (1) | 0.499 |
| Possible | 0.4 (3) | 0.3 (2) | 1.000 |
| Definite/probable | 0.1 (1) | 0.8 (6) | 0.069 |
| Any | 0.5 (4) | 1.1 (8) | 0.264 |
| Cumulative (0–720 days) | | | |
| Definite | 0.8 (6) | 0.8 (6) | 1.000 |
| Probable | 0.3 (2) | 0.1 (1) | 1.000 |
| Possible | 0.8 (6) | 0.7 (5) | 1.000 |
| Definite/probable | 1.1 (8) | 0.9 (7) | 1.000 |
| Any | 1.9 (14) | 1.6 (12) | 0.844 |

Values are % (n).
Abbreviations as in Table 1.

| Table 4. ARC Definite/Probable Stent Thrombosis Up to 2 Years | | | | |
|---|----------------|----------------------|----------------|-------------------|
| DES | ARC Definition | Time to Event (Days) | Clinical Event | DAPT at Event |
| PES | Definite | 413 | Q-wave MI | ASA only |
| | Definite | 495 | Non-Q-wave MI | ASA + clopidogrel |
| | Definite | 619 | Non-Q-wave MI | ASA + clopidogrel |
| | Definite | 645 | Q-wave MI | ASA only |
| | Definite | 689 | Non-Q-wave MI | ASA only |
| | Probable | 697 | Q-wave MI | ASA + clopidogrel |
| ZES | Definite | 369 | Q-wave MI | None |


ARC = Academic Research Consortium; DAPT = dual antiplatelet therapy; DES = drug-eluting stent(s); other abbreviations as in Table 1.

Q-wave and 3 non-Q-wave MIs, and 3 of the 6 patients were no longer receiving clopidogrel therapy (Table 4). The sole ZES VLST case was associated with a Q-wave MI in a patient who stopped taking both aspirin and clopidogrel 10 days before the ST event. The cumulative incidence of ARC definite or probable ST after 1 year mirrored the cumulative incidence of MIs after 1 year (Fig. 5). **Cost-effectiveness.** There were no differences in quality-adjusted survival or medical costs (initially, during the second year, or cumulative during the 2 years of follow-up) between ZES and PES patient cohorts (Table 5). Similarly, bootstrap analysis of the 2-year cost-effectiveness showed no benefit of ZES compared with PES patients (Fig. 6).

Discussion

The key findings of the present analysis assessing clinical outcomes after 2 years in patients treated with ZES versus those treated with PES for single de novo coronary lesions from the randomized ENDEAVOR IV trial (1,548 patients with >95% follow-up) are: 1) up to 2 years after stent

implantation, the ZES—compared with the PES—continued to demonstrate comparable clinical anti-restenosis efficacy; 2) ZES seemed to have a more favorable mid-term safety profile, with fewer episodes of VLST and significantly less MI in patients compared with PES; and 3) at 2 years, there was similar cost-effectiveness comparing ZES and PES. Very late stent thrombosis and in-stent restenosis, the main 2 concerns after DES implantation, require meticulous long-term clinical follow-up to ensure patient safety and to determine whether overall clinical effectiveness is durable (7–9). Despite significantly higher 8-month angiographic in-stent late loss in ZES-treated patients, ischemia-driven TLR between 1 and 2 years were infrequent in the ENDEAVOR IV trial and similar for both ZES (1.6%) and PES (1.7%), and cumulative ischemia-driven TLR at 2 years was likewise similar for ZES (5.9%) and PES (4.6%). The clinical efficacy results for the ENDEAVOR IV trial at 2 years are consistent with multiple other ZES studies (35–39), suggesting the absence of a “late catch-up” phe-






Figure 5. Landmark Analysis of Incidence of Myocardial Infarction During the First Year and Between the First and Second Years After ZES and PES Implantation

The HRs with 95% confidence intervals for 0 to 1 year and 1 to 2 years are shown for the incidence rates of myocardial infarction. Abbreviations as in Figures 1 and 2.

nomenon and a durable effect on the reduction of clinical restenosis over time (26,38,40–42). Because recent reports have indicated concerns associated with increasing late loss and related late TLR events after 1 year with other DES (43,44), these ZES findings bear watching as further clinical follow-up is accumulated in the ENDEAVOR IV trial and other ZES clinical trials involving more complex lesion subsets (39,45). It is interesting to note that the 2-year results from the ENDEAVOR IV trial continue to emphasize the significant impact of angiographic follow-up on subsequent TLR events. In the subset of patients with planned angiographic follow-up, there were significant differences in 2-year TLR rates (ZES 9.2% vs. PES 3.1%, $p = 0.045$), whereas in those patients without angiographic follow-up (82% of the study population), there were no differences in 2-year TLR (ZES 5.2% vs. PES 4.9%, $p = 0.896$).

Because stent-related coronary thrombosis during the first year after implantation is known to be associated with

multiple patient, lesion, and procedural predictors (25,46,47) that vary from study to study, we have emphasized the importance of very late (after 1 year) thrombosis events, which seem to be related to fundamental design characteristics of DES. The 2-year clinical outcomes from the ENDEAVOR IV trial reinforce the impression that ZES are similar to BMS from the standpoint of late safety, including VLST. However, it bears noting that the ENDEAVOR IV trial was not powered to demonstrate differences in safety end points such as cardiac death plus MI or ST, and after 2-year follow-up, the only statistically significant benefit of ZES over PES was a reduction in non-Q-wave MIs (3.5% vs. 1.6%, $p = 0.022$). Moreover, there are data from other recent randomized trials similarly underpowered to discern differences in ST, which compare ZES with other DES (both SES and PES) during the first year of follow-up. These studies show no consistent differences in ST during the first year of clinical follow-up (48,49). The largest randomized trial (8,800 patients, with enrollment

Table 5. 2-Year QA Survival and Medical Costs

| | Taxus | Endeavor | Difference (95% CI) | p Value |
|----------------------------------|--------|----------|-----------------------|---------|
| Survival (days) | 711.22 | 711.00 | 0.21 (–6.07 to 6.49) | 0.95 |
| Discount survival (days)* | 690.09 | 689.89 | 0.20 (–5.85 to 6.24) | 0.95 |
| Quality-adjusted survival (days) | 575.41 | 576.00 | –0.60 (–5.95 to 4.75) | 0.83 |
| Discount QA survival (days)* | 557.97 | 558.56 | –0.59 (–5.74 to 4.56) | 0.82 |
| Medical costs (\$) | | | | |
| Initial year | 17,713 | 17,167 | 545 (–178 to 1,269) | 0.14 |
| Second year | 2,316 | 2,454 | –137 (–782 to 510) | 0.68 |
| Cumulative 2-yr costs | 20,029 | 19,621 | 408 (–632 to 1,269) | 0.44 |
| Cumulative discount 2-yr costs* | 19,852 | 19,440 | 411 (–600 to 1,423) | 0.43 |

*Discount rate of 3%/year.

CI = confidence interval; QA = quality-adjusted.




Figure 6. Bootstrap Results for the 2-Year Cost-Effectiveness

Results are shown as discounted mean values for the cost-effectiveness ratio.

completed) focusing on the comparative safety of ZES versus SES was designed to provide a more definitive answer to long-term (3-year) safety in a real-world patient population (45).

In the ENDEAVOR IV trial, there were more MIs in the PES group after 1 year due to a higher incidence of VLST events (1 event with ZES vs. 6 with PES). The 1 ZES patient with VLST had stopped taking both ASA and clopidogrel 10 days before the thrombosis event. These findings are similar to previous ZES studies versus BMS or SES (37,40,41). Moreover, a pooled analysis of all adjudicated ZES clinical trials with long-term follow-up indicated a VLST frequency of only 0.2% in over 2,000 ZES patients (26,42). Combined with higher rates of periprocedural MIs in PES patients, possibly related to increased side branch occlusions associated with PES stents (5,50), the overall early and late safety profile of ZES in the ENDEAVOR IV trial was favorable compared with PES.

The comprehensive cost-effectiveness analysis that was imbedded in the ENDEAVOR IV study design indicated no differences in quality-adjusted survival, medical costs, and cost-effectiveness, either acutely or at 2 years, comparing ZES versus PES treatment.

Study limitations. The patients and lesions treated in the ENDEAVOR IV study were not highly complex, and therefore, the results of this study cannot be extrapolated to patients with more complex coronary anatomy. This study was not designed or powered to compare rates of infrequent adverse clinical events such as death, MI, or ST; therefore,

the results and trends should be interpreted as largely hypothesis-generating. Finally, longer follow-up is necessary to adequately assess late safety and efficacy events when comparing ZES and PES therapies.

Conclusions

This analysis of the ENDEAVOR IV trial demonstrates that the initial comparable clinical efficacy of ZES compared with PES is sustained at 2 years, with a similarly low rate of TLR and no differences in either TVF or MACE. Comparable clinical outcomes resulted in similar 2-year cost-effectiveness. These findings, coupled with the low incidence of VLST and lower incidence of MI after ZES versus PES implantation, provide evidence supporting the mid-term safety and efficacy of ZES in the treatment of de novo lesions in native coronary arteries. Longer follow-up of the ENDEAVOR IV trial and confirmation from additional randomized trials are necessary to substantiate these findings.

Acknowledgments

The authors would like to thank Naureen Sheikh, PhD, and Denise Jones, BSN, Medtronic CardioVascular, for management of study clinical operations. The authors would like to acknowledge the important contributions of Manuela Negoita, MD, Minglei Liu, PhD, and Colleen Gilbert, PharmD, with manuscript preparation. Similarly, Dr. Leroy LeNarz was an invaluable collaborator in the design, exe-

cution, and analysis of results from the ENDEAVOR IV trial.

Reprint requests and correspondence: Dr. Martin B. Leon, Professor of Medicine and Associate Director, Center for Interventional Vascular Therapy, Columbia University Medical Center, New York, New York 10032. E-mail: mleon@crf.org.

REFERENCES

1. Moses JW, Leon MB, Popma JJ, et al. Sirolimus-eluting stents versus standard stents in patients with stenosis in a native coronary artery. *N Engl J Med* 2003;349:1315-23.
2. Stone GW, Ellis SG, Cox DA, et al. A polymer-based, paclitaxel-eluting stent in patients with coronary artery disease. *N Engl J Med* 2004;350:221-31.
3. Stone GW, Ellis SG, Cannon L, et al. Comparison of a polymer-based paclitaxel-eluting stent with a bare metal stent in patients with complex coronary artery disease: a randomized controlled trial. *JAMA* 2005;294:1215-23.
4. Dawkins KD, Grube E, Guagliumi G, et al. Clinical efficacy of polymer-based paclitaxel-eluting stents in the treatment of complex, long coronary artery lesions from a multicenter, randomized trial: support for the use of drug-eluting stents in contemporary clinical practice. *Circulation* 2005;112:3306-13.
5. Leon MB, Mauri L, Popma JJ, et al. A randomized comparison of the ENDEAVOR zotarolimus-eluting stent versus the TAXUS paclitaxel-eluting stent in de novo native coronary lesions: 12-month outcomes from the ENDEAVOR IV trial. *J Am Coll Cardiol* 2009. In press.
6. Stone GW, Midei M, Newman W, et al. Comparison of an everolimus-eluting stent and a paclitaxel-eluting stent in patients with coronary artery disease. A randomized trial. *JAMA* 2008;299:1903-13.
7. Camenzind E, Steg PG, Wijns W. Stent thrombosis late after implantation of first-generation drug-eluting stents: a cause for concern. *Circulation* 2007;115:1440-55.
8. Daemen J, Wenaweser P, Tsuchida K, et al. Early and late coronary stent thrombosis of sirolimus-eluting and paclitaxel-eluting stents in routine clinical practice: data from a large two-institutional cohort study. *Lancet* 2007;369:667-78.
9. Lagerqvist B, James SK, Stenestrand U, et al. Long-term outcomes with drug-eluting stents versus bare-metal stents in Sweden. *N Engl J Med* 2007;356:1009-19.
10. Machara A, Mintz GS, Weissman NJ, et al. Late thrombosis after gamma-brachytherapy. *Catheter Cardiovasc Interv* 2003;58:455-8.
11. Waksman R, Ajani AE, Pinnow E, et al. Twelve versus six months of clopidogrel to reduce major cardiac events in patients undergoing gamma-radiation therapy for in-stent restenosis: Washington Radiation for In-Stent restenosis Trial (WRIST) 12 versus WRIST PLUS. *Circulation* 2002;106:776-8.
12. Wenaweser P, Daemen J, Zwahlen M, et al. Incidence and correlates of drug-eluting stent thrombosis in routine clinical practice. 4-year results from a large 2-institutional cohort study. *J Am Coll Cardiol* 2008;52:1134-40.
13. Ferrari E, Benhamou M, Cerboni P, Marcel B. Coronary syndromes following aspirin withdrawal: a special risk for late stent thrombosis. *J Am Coll Cardiol* 2005;45:456-9.
14. Airolidi F, Colombo A, Morici N, et al. Incidence and predictors of drug-eluting stent thrombosis during and after discontinuation of thienopyridine treatment. *Circulation* 2007;116:745-54.
15. Spertus JA, Kettelkamp R, Vance C, et al. Prevalence, predictors, and outcomes of premature discontinuation of thienopyridine therapy after drug-eluting stent placement: results from the PREMIER registry. *Circulation* 2006;113:2803-9.
16. Grines CL, Bonow RO, Casey DE Jr., et al., American Heart Association; American College of Cardiology; Society for Cardiovascular Angiography and Interventions; American College of Surgeons; American Dental Association; American College of Physicians. Prevention of premature discontinuation of dual antiplatelet therapy in patients with coronary artery stents: a science advisory from the American Heart Association, American College of Cardiology, Society for Cardiovascular Angiography and Interventions, American College of Surgeons, and American Dental Association, with representation from the American College of Physicians. *J Am Coll Cardiol* 2007;49:734-9.
17. Hodgson JM, Stone GW, Lincoff AM, et al., Society for Cardiovascular Angiography and Interventions. Late stent thrombosis: considerations and practical advice for the use of drug-eluting stents: a report from the Society for Cardiovascular Angiography and Interventions Drug-eluting Stent Task Force. *Catheter Cardiovasc Interv* 2007;69:327-33.
18. Joner M, Finn AV, Farb A, et al. Pathology of drug-eluting stents in humans: delayed healing and late thrombotic risk. *J Am Coll Cardiol* 2006;48:193-202.
19. Finn AV, Joner M, Nakazawa G, et al. Pathological correlates of late drug-eluting stent thrombosis: strut coverage as a marker of endothelialization. *Circulation* 2007;115:2435-41.
20. Joner M, Nakazawa G, Finn AV, et al. Endothelial cell recovery between comparator polymer-based drug-eluting stents. *J Am Coll Cardiol* 2008;52:333-42.
21. Awata M, Nanto S, Uematsu M, et al. Angioscopic comparison of neointimal coverage between zotarolimus- and sirolimus-eluting stents. *J Am Coll Cardiol* 2008;52:789-90.
22. Guagliumi G, Musumeci G, Sirbu V, et al. A prospective, randomized, controlled study using optical coherence tomography to evaluate strut coverage of sirolimus-, paclitaxel-, and zotarolimus-eluting coronary stents in long lesions requiring overlapping. Paper presented at: Transcatheter Cardiovascular Therapeutics 2008, October 14, 2008; Washington, DC.
23. Shin DI, Seung KB, Kim PJ, et al. Long-term coronary endothelial function after zotarolimus-eluting stent implantation. *Int Heart J* 2008;49:639-52.
24. Kim JW, Seo HS, Park JH, et al. A prospective, randomized, 6-month comparison of the coronary vasomotor response associated with a zotarolimus- versus a sirolimus-eluting stent: differential recovery of coronary endothelial dysfunction. *J Am Coll Cardiol* 2009;53:1653-9.
25. Cutlip DE, Baim DS, Ho KK, et al. Stent thrombosis in the modern era: a pooled analysis of multicenter coronary stent clinical trials. *Circulation* 2001;103:1967-71.
26. Mauri L, Cutlip DE, Leon MB, et al. Long term safety results from the Endeavor program: four-year follow-up. *Am J Cardiol* 2008;102 Suppl:124i.
27. Cutlip DE, Windecker S, Mehran R, et al., on behalf of the Academic Research Consortium. Clinical end points in coronary stent trials: a case for standardized definitions. *Circulation* 2007;115:2344-51.
28. The Trees. Version 25.0. 2008. CMS MS-DRG Grouper. Eden Prairie, MN: Ingenix Corporation. Available at: www.ingenix.com. Accessed June 1, 2009.
29. DRG Expert: A Comprehensive Guidebook to the DRG Classification System. Salt Lake City, UT: Ingenix, 2008.
30. DRG Plus! 2008: A Comprehensive Guide to Medicare Severity Diagnosis Related Groups. Los Angeles, CA: Practice Management Information Corporation, 2007.
31. Mitchel J, Burge R, Lee A, et al. Per Case Prospective Payment for Episodes of Hospital Care. PB95226023. 1995. Springfield, VA: U.S. Department of Commerce National Technical Information Service.
32. Konstane RP, Eisenstein EL, Anstrom KJ, et al. Outcomes of second revascularization procedures after stent implantation. *J Med Syst* 2008;32:177-86.
33. Garg P, Cohen DJ, Gaziano T, Mauri L. Balancing the risks of restenosis and stent thrombosis in bare-metal versus drug-eluting stents: results of a decision analytic model. *J Am Coll Cardiol* 2008;51:1844-53.
34. Tsevat J, Goldman L, Soukup JR, et al. Stability of time-tradeoff utilities in survivors of myocardial infarction. *Med Decis Making* 1993;13:161-5.
35. Meredith IT, Ormiston J, Whitbourn R, et al. First-in-human study of the Endeavor ABT-578-eluting phosphorylcholine-encapsulated stent system in de novo native coronary artery lesions: ENDEAVOR I trial. *EuroIntervention* 2005;1:157-64.

36. Fajadet J, Wijns W, Laarman GJ, et al., ENDEAVOR II Investigators. Randomized, double-blind, multicenter study of the Endeavor zotarolimus-eluting phosphorylcholine-encapsulated stent for treatment of native coronary artery lesions: clinical and angiographic results of the ENDEAVOR II trial. *Circulation* 2006;114:798-806.
37. Kandzari DE, Leon MB, Popma JJ, et al., ENDEAVOR III Investigators. Comparison of zotarolimus-eluting and sirolimus-eluting stents in patients with native coronary artery disease: a randomized controlled trial. *J Am Coll Cardiol* 2006;48:2440-7.
38. Gershlick A, Kandzari DE, Leon MB, et al., ENDEAVOR Investigators. Zotarolimus-eluting stents in patients with native coronary artery disease: clinical and angiographic outcomes in 1,317 patients. *Am J Cardiol* 2007;100:45M-55M.
39. Jain AK, Meredith IT, Lotan C, Rothman MT, Pateraki S, E-Five Investigators. Real-world safety and efficacy of the endeavor zotarolimus-eluting stent: early data from the E-Five Registry. *Am J Cardiol* 2007;100:77M-83M.
40. Meredith IT, Ormiston J, Whitbourn R, et al., ENDEAVOR I Investigators. Four-year clinical follow-up after implantation of the endeavor zotarolimus-eluting stent: ENDEAVOR I, the first-in-human study. *Am J Cardiol* 2007;100:56M-61M.
41. Wijns W, Fajadet J, Laarman GM, et al. Randomized trial comparing the Endeavor zotarolimus-eluting stent and the Driver bare-metal stent in single de novo native coronary artery lesions: five-year clinical follow-up of ENDEAVOR II. *EuroIntervention* 2009;5 Suppl E:E27.
42. Kandzari DE. Long term efficacy outcomes following treatment with zotarolimus-eluting stents: analysis of the pooled Endeavor program. Presented at: EuroPCR; May 19, 2009; Barcelona, Spain.
43. Byrne RA, Iijima R, Mehilli J, et al. Durability of antirestenotic efficacy in drug-eluting stents with and without permanent polymer. *J Am Coll Cardiol Interv* 2009;2:291-9.
44. Finn AV, Nakazawa G, Kolodgie FD, Virmani R. Temporal course of neointimal formation after drug-eluting stent placement: is our understanding of restenosis changing? *J Am Coll Cardiol Interv* 2009;2:300-2.
45. Brugaletta S, Burzotta F, Sabaté M. Zotarolimus for the treatment of coronary artery disease: pathophysiology, DES design, clinical evaluation and future perspective. *Expert Opin Pharmacother* 2009;10:1047-58.
46. Hausleiter J, Kastrati A, Mehilli J, et al. Predictive factors for early cardiac events and angiographic restenosis after coronary stent placement in small coronary arteries. *J Am Coll Cardiol* 2002;40:882-9.
47. Uren NG, Schwarzacher SP, Metz JA, et al., POST Registry Investigators. Predictors and outcomes of stent thrombosis: an intravascular ultrasound registry. *Eur Heart J* 2002;23:124-32.
48. Lassen JF, Rasmussen K, Galløe A, et al. SORT-OUT III: A Prospective, Randomized Comparison of Zotarolimus-eluting and Sirolimus-eluting Stents in Patients with Coronary Artery Disease. Paper presented at: Transcatheter Therapeutics; October 16, 2008; Washington, DC.
49. Park SJ. The Zotarolimus-Eluting Stent Versus Sirolimus-Eluting Stent and Paclitaxel-Eluting Stent for Coronary Lesions (ZEST) trial. Paper presented at: i2 Summit at ACC; March 29, 2009; Orlando, FL.
50. Popma JJ, Mauri L, Kirtane A, Cutlip D, Almonacid A, Leon MB. Frequency and clinical consequences associated with sidebranch occlusion during stent implantation using zotarolimus-eluting and paclitaxel-eluting coronary stents. *Circ Cardiovasc Intervent* 2009;2:133-9.

Key Words: angioplasty ■ coronary disease ■ drug-eluting stent ■ revascularization ■ stent thrombosis.